




A JOURNAL ON TAXONOMIC BOTANY,
PLANT SOCIOLOGY AND ECOLOGY



REINWARDTIA

12(1)

REINWARDTIA

*A JOURNAL ON TAXONOMIC BOTANY,
PLANT SOCIOLOGY AND ECOLOGY*

Vol. 12(1): 1-128.22 July 2002

Editors

ELIZABETH A. WIDJAJA, MIEN A. RIFAI, SOEDARSONO RISWAN, JOHANIS P. MOGEA

Correspondence and subscriptions of the journal should be addressed to

HERBARIUM BOGORIENSE, BIDANG BOTANI, PUSAT PENELITIAN BIOLOGI - LIPI,
BOGOR, INDONESIA

A NEW SPECIES OF DALBERGIA (LEGUMINOSAE) FROM MALAY PENINSULA

BAMBANG SUNARNO

Herbarium Bogoriense, Bidang Botani, Puslit Biologi-LIPI, Indonesia

& HIROSHI OHASHI

Biological Institute, Graduate School of Science, Tohoku University, Sendai, Japan

ABSTRACT

SUNARNO, BAMBANG & OHASHI, HIROSHI. 2002. A new species of *Dalbergia* (Leguminosae) from Malay Peninsula. *Reinwardtia* 12(1): 117–119. — A new species, *Dalbergia johorensis* from the Malay Peninsula is described. It is close to *D. rostrata* and *D. havilandii* but readily distinguished by the grooved midrib beneath, flowers with narrower standard and wings and style hairy in the lower part.

Keywords: *Dalbergia*, *Leguminosae*, Malay Peninsula.

ABSTRAK

SUNARNO, BAMBANG & OHASHI, HIROSHI. 2002. Jenis baru *Dalbergia* (Leguminosae) dari Semenanjung Malaya. *Reinwardtia* 12(1): 117–119. — Dipertelakan suatu jenis baru *Dalbergia johorensis* dari Semenanjung Malaya. Jenis ini dekat dengan *D. rostrata* dan *D. havilandii* namun dapat dibedakan dari permukaan bawah tulang daun yang beralur, bunga dengan bendera dan sayap yang lebih menyempit serta bagian bawah tangkai putiknya berbulu.

Kata kunci: *Dalbergia*, *Leguminosae*, Semenanjung Malaya

For a revision of the genus *Dalbergia* in Malesia we have published some new species from Sulawesi and Borneo (Sunarno & Ohashi, 1996; 1997). This paper is merely a continuation of the previous work for the Malesian *Dalbergia*. During our examination to the specimens in Herbarium Bogoriense (BO) Indonesia and Rijksherbarium (L) the Netherlands, we identified about 11 species are occurred in the Peninsular Malaysia, one of which is unknown to science. Similar specimen as found from Borneo was referred to as *Dalbergia* sp. due to its incompleteness (Sunarno & Ohashi, 1997). The finding of complete specimen from Johore lead us to consider it to be a new species. The species is similar to *D. rostrata* Hassk., a species widely distributed in Malesia except New Guinea, and to *D. havilandii* Prain, a bornean species (Prain 1904) but they differ in leaves, stipules and flowers (Table 1). In this paper the new species is described and illustrated.

***Dalbergia johorensis* Sunarno et Ohashi, sp. nov.**
– Fig. 1.

Dalbergia foliolis rostratis sed differt floris vexillae et alae angustiora (ca. 1.5 mm longa), stylus

sparsus pubescentibus infra parte, infra costa sulco in longitudinem. – Typus: Malay Peninsula, Johore, Namheng estate, Kota Tinggi, fl. Febr., 1930. (*Teruya* 1192 BO–Holo; SING– Iso).

Liana. Branchlets stoutish, terete, sparsely or rather densely brown puberulent, striated downwards, lenticels gradually inconspicuous. Stipules very early caducous, ovate, ca. 2 x 1 mm, densely puberulent. Leaves 2.0–4.5 cm long; petioles 1.5–2.5 cm long, densely brown puberulent; rachis 0.5–2 cm long, densely rusty puberulent. Leaflets 1–3, oblong or ovate, rounded or slightly cordate base, obtuse, apiculate and acutely acuminate towards the very apex, 5–7 x 3–4 cm, firmly coriaceous, light brown and glabrous above, finely reticulate, densely adpressed short brown puberulent below; the midrib distinctly grooved above and prominently so beneath, sparsely to rather densely brown puberulent; secondary nerves 7–9 pairs; margin slightly prominently revolute; petioles 0.4–0.5 cm long, densely puberulent but gradually sparsely so downwards; bracts caducous, ovate, 2–3 x 1–2 mm, outer surface densely rusty puberulent. Inflorescences axillary, panicles of raceme, up to 20 cm long, peduncles up to 9 cm long, sparsely puberulent;

Table 1. Comparison of *D. havilandii*, *D. johoriensis* and *D. rostrata*

| Morphological Characters | <i>D. havilandii</i> | <i>D. johoriensis</i> | <i>D. rostrata</i> |
|----------------------------------|---|---|---|
| Stipules | subpersistent, ovate-lanceolate, 6.5—7 mm long | caducous, ovate, ca. 2 mm long | caducous, broadly ovate, ca. 1 mm long |
| Leaflets number | 1—3 | 1—3 | 1—5 |
| Size | 5—7.5 x 2.5—4 cm | 5—7 x 3—4 cm | 4—5 x 2—9 cm |
| Base | obtuse to truncate | rounded or slightly cordate | cuneate, rounded or obtuse |
| Apex | obtuse or acute | rostrate, apiculate | rostrate, slightly cuspidate |
| Surface | densely hairy on both surfaces | sparsely hairy on both surfaces | glabrous upper surface, sparsely hairy beneath |
| Midrib beneath of lateral nerves | not grooved 4—5 pairs | grooved 7—9 pairs | not grooved 7—9 pairs |
| Inflorescence | short clustered raceme, many flowers | panicle of racemes few flowers | panicle of raceme or compound panicle, many flowers |
| Bracteoles | subpersistent narrowly ovate or linear, ca. 1 mm long | persistent, narrowly ovate or linear, ca. 1 mm long | caducous, narrowly ovate or linear, ca. 1 mm long |
| Flowers size | Unknown | 6—7(-8) mm long | 6—7 mm long |
| Calyx | 3—3.5 mm long | ca. 4 mm long | 3.5—4.5 mm long |
| Standard | orbicular, apex rounded, emarginate | broadly obovate, apex rounded, entire | suborbicular, apex rounded, emarginated |
| Pistil | style glabrous | style sparsely hairy at lower point | style glabrous |

rachis ca. 10 cm long, sparsely puberulent but gradually sparsely so downwards; bracts caducous, ovate, 2–3 x 1–2 mm, outer surface densely rusty puberulent, bracteoles persistent, linear or narrowly ovate, ca. 1 mm long, densely rusty puberulent. Flowers 6–7 (–8) mm long, pedicels stoutish, 1–2 mm long, densely brown puberulent. Calyx bell-shaped, densely rusty pubescent, tube ca. 3 mm long; lobes subequal, the lowest longer than others, ca. 1 mm long. Standards reflexed, blades broadly obovate, 2.5–3 x ca. 2.5 mm, attenuate to base, rounded, entire at apex, finely reticulated; claw 2.5–3 mm long. Wing blades oblong, ca. 2.5 x ca. 1.5 mm; base acute or obtuse one side, rounded on the other; apex obtuse to rounded; claw ca. 3 mm long. Keel blades boat-shaped, ca. 2.5 x 1.5 mm; base obtuse on one side, rounded on the other, united in the upper side; claw ca. 3 mm long. Stamens 9, monadelphous, the united filaments 3.5–4 mm long, the upper free one 1–1.5(–2) mm long. Pistil 5.5–6.0 mm long, ovary densely hairy; stipe ca. 2 mm long, glabrous; style ca. 1.5 mm long, sparsely rusty puberulent in the lower part; ovules 1–2. Pods unknown.

DISTRIBUTION. Malay Peninsula (Johore) and Borneo (Sarawak; Kuching).




Fig.1. *Dalbergia johorensis* Sunarno et H. Ohashi. A. Branch with unifoliate and trifoliate leaves. B. Inflorescences. C. Part of leaflet showing a groove along the midrib beneath. D. Flowers. E. Petals. F. Calyx (opened). G. Stamens. H. Pistil (from Teruya 1192).

HABITAT AND ECOLOGY. Roadsides in thickets. Flowering in February and September.

NOTES. The number of the leaflets is close to *D. rostrata* and *D. havilandii*, however, comparison to those species (Table 1) indicates that the current species is different.

SPECIMENS EXAMINED. Malay Peninsula, Johore, Namheng estate, Kota Tinggi, fl. Febr. 1930. *Teruya* 1192. (BO-Holo.).

ACKNOWLEDGEMENTS

The first author would like to express his gratitude to the British Council and Flora Malesiana Foundation for financial support during his studies in Royal Botanical Gardens, Kew, and Rijksherbarium, Leiden. He is grateful to Drs. R.M. Polhill and B. Verdcourt of Kew, Drs. P. Baas, the late C. Kalkman, the late R.

Geesink for their support and encouragement for the study. We wish to thank Drs. Nemoto of Biological Institute of Tohoku University and H. Wiriadinata of Herbarium Bogoriense for their advice and help. This work was accomplished as a part of doctoral dissertation submitted by the first author to Tohoku University, Sendai, with financial supports to us from Japan Society for the promotion of Science (JSPS) in 1995–1996.

REFERENCES

- PRAIN, D. 1904. The species of *Dalbergia* of Southeastern Asia. *Ann. Roy. Bot. Gard. Calc.* 10: 1–114, pls. 1–91.
- SUNARNO, B. & OHASHI, H. 1996. *Dalbergia* (*Leguminosae*) of Sulawesi, Indonesia. *Journ. Jap. Bot.* 71 (5): 241–248.
- SUNARNO, B. & OHASHI, H. 1997. *Dalbergia* (*Leguminosae*) of Borneo. *Journ. Jap. Bot.* 72 (4): 198–220.

INSTRUCTION TO AUTHORS

Manuscripts intended for publication in *Reinwardtia* should be written either in English, French or German, and represent articles which have not been published in any other journal or proceedings. Each manuscript received will be considered and processed further if it is accompanied by signed statements given independently by two reviewers chosen by the author(s) attesting to its merits as well as its scientific suitability for publication in *Reinwardtia*.

Two printed copies (on A4 paper) of the manuscript of not more than 200 pages should be sent to Editors, together with an electronic copy prepared on Word Processor computer programme using Times New Romance letter type and saved as Rich Text File must be submitted.

For the style of presentation authors should follow the latest issue of *Reinwardtia* very closely. Title of the article should be followed by author's name and mailing address and a one-paragraphed abstract in English (with French or German abstract for papers in French or German) of not more than 250 words. Keywords should be given below each abstract. On a separate paper author(s) should prepare the preferred running title of the article submitted.

Taxonomic keys should be prepared using the aligned-couplet type.

Strict adherence to the *International Code of Botanical Nomenclature* is observed, so that taxonomic and nomenclatural novelties should be clearly shown, Latin description for new taxon proposed should be provided, and the herbaria where type specimens are deposited should be indicated. Synonyms should be presented in the long form [name of taxon, author's name, year of publication, abbreviated journal or book title, volume (number): page].

Maps, line drawing illustrations or photographs preferably should be prepared in landscape presentation to occupy two columns. Illustrations must be submitted as original art accompanying, but separate from, the manuscripts. On electronic copy, the illustrations should be saved in jpg or .gif format. Legends for illustrations must be submitted separately at the end of the manuscript.

Bibliography, list of literature cited or references follow the Harvard System.

For each paper published author(s) will receive 25 copies of reprints free of charge. Any additional copies should be ordered in advance and the author(s) will be charged accordingly.

ISSN 0034 - 365 X

REINWARDTIA

Vol. 12. No. 1.2002

CONTENTS

Page

Y. PURWANTO. Gestion de la Biodiversite: Relations aux plantes et dynamiques
vegetales chez les Dani de la Vallee de la Baliem en Irian Jaya, Indonesie.1

TRI MULYANINGSIH & COLIN ERNEST RIDSDALE. The Bornean genus *Hypo-*
bathrum (Rubiaceae): an investigation of its characters and taxonomic status.95

BAMBANG SUNARNO & HIROSHI OHASHI. A new species of *Dalbergia* (*Legumi-*
nosae) from Malay Peninsula.117

BAMBANG SUNARNO. New species of *Labisia* (*Myrsinaceae*) from Sumatra.121

SRI S. TJITROSOEDIRDJO. Four new taxa of *Asteraceae* in Sumatra.125